

Localised edge states nucleate turbulence in extended plane Couette cells

Tobias M. Schneider ^{a,b,*}; Daniel Marinc ^{a,c,†}
and Bruno Eckhardt ^{a,d,‡}

^aFachbereich Physik, Philipps-Universität Marburg,
D-35032 Marburg, Germany

^bSchool of Engineering and Applied Sciences,
Harvard University, Cambridge, MA 02138, USA

^cAerodynamisches Institut, RWTH Aachen, D-52062 Aachen,
Germany

^dDepartment of Mechanical Engineering, TU Delft,
2628 CA Delft, The Netherlands

September 3, 2009

Abstract

We study the turbulence transition of plane Couette flow in large domains where localised perturbations are observed to generate growing turbulent spots. Extending previous studies on the boundary between laminar and turbulent dynamics we determine invariant structures intermediate between laminar and turbulent flow. In wide but short domains we find states that are localised in spanwise direction, and in wide and long domains the states are also localised in downstream direction. These localised states act as critical nuclei for the transition to turbulence in spatially extended domains.

1 Introduction

Landau (1944) is accredited with a description of the transition to turbulence through a sequence of instabilities that add spatial and temporal degrees of freedom to the flow, thereby building up the complex dynamics which characterise turbulent motion. Such a scenario seems to be realised in fluids heated from below (Rayleigh-Bénard) or in centrifugally unstable situations (Taylor-Couette)

*tschneid@seas.harvard.edu

†d.marinc@aia.rwth-aachen.de

‡bruno.eckhardt@physik.uni-marburg.de

(Koschmieder, 1993), but it does not apply to several important flows like pressure driven flow down a circular pipe (Reynolds, 1883) or plane Couette flow (Schmid & Henningson, 1999; Eckhardt *et al.*, 2007), since already the first step of the classical scenario, the linear instability of the laminar profile, is missing. Accordingly, triggering turbulence in these linearly stable flows requires that both the flow rate *and* the strength of an applied perturbation exceed critical levels (Boberg & Brosa, 1988; Grossmann, 2000). Several experimental (Darbyshire & Mullin, 1995; Hof *et al.*, 2003; Peixinho & Mullin, 2007; Dauchot & Daviaud, 1995; Bottin *et al.*, 1998*a,b*) and numerical studies (Meseguer, 2003; Schneider *et al.*, 2007; Schmiegel & Eckhardt, 1997; Eckhardt *et al.*, 2008) have focused on the required minimal perturbations and have identified a very sensitive dependence of the critical amplitudes on both the spatial structure of a perturbation and on the flow rate.

From a dynamical systems point of view, the coexistence of the stable laminar profile with turbulent dynamics implies that there is a boundary in the state space of the system which separates perturbations that return to the laminar profile from those that become turbulent (Eckhardt *et al.*, 2002, 2007). The sensitive dependence on initial conditions results in a fractal and convoluted boundary which was termed *edge of chaos* (Skufca *et al.*, 2006; Schneider *et al.*, 2007; Vollmer *et al.*, 2009). It generalises the more familiar basin boundary to situations where turbulence might be transient (Hof *et al.*, 2006). Despite its intricate geometry the edge is locally formed by the stable set of an invariant dynamical object called *edge state*. By definition, the edge state corresponds to a self-sustained non-laminar and non-decaying flow field of critical energy. Its stable manifold is of co-dimension one per construction and defines locally the stability boundary. Therefore, the edge state together with its stable manifold determines minimal seeds for turbulence.

Both the exact solutions that have been linked to the turbulent dynamics (Nagata, 1990; Clever & Busse, 1997; Waleffe, 2003; Eckhardt *et al.*, 2008) and the edge states (Wang *et al.*, 2007; Schneider *et al.*, 2008) which guide the transition have been studied in small computational domains subject to periodic boundary conditions. Thus, they focus on the temporal degrees of freedom but cannot capture large scale spatial phenomena such as the growth of turbulent regions or the coexistence of turbulent and non-turbulent patterns observed in spatially extended systems. The spatial dynamics of extended flow systems shows up in transition experiments where the homogeneously driven flow is locally perturbed by a jet injection (Bottin *et al.*, 1998*b*) or a small obstacle (Bottin *et al.*, 1997): in such cases one first observes a localised turbulent region which then starts to spread out (Emmons, 1951). The spatially extended edge states cannot explain these phenomena, since they would require that the perturbation exceeds the critical threshold everywhere in space, in contrast to the experimental evidence. Studies on pipe flow, both in a model (Duguet *et al.*, 2008) and in the fully resolved direct numerical simulations (Mellibovsky *et al.*, 2009) have identified an edge state that is localised along the axis. In the present study we apply these ideas and tools to the case of plane Couette flow, where there are two directions of spatial extension, streamwise and span-

wise. Using direct numerical simulations we show that these edge states can be localised in one or both directions, thereby confirming the expectation that a localised perturbation should be sufficient to nucleate turbulence. Moreover, we find tantalising similarities to observations in typical pattern forming systems (Knobloch, 2008).

2 Edge states in wide Couette systems

As usual, we define the Reynolds number for plane Couette flow as $Re = u_0 d / \nu$, where u_0 is half the velocity differences between the two plates, d is half the gap width d and ν the viscosity of the fluid. In the following all lengths will be given in units of d . The system is transitionally invariant in both the streamwise (x) and spanwise (z) direction. The laminar linear flow profile is stable against infinitesimal perturbations for all Re (Schmid & Henningson, 1999). In the turbulent case the translational symmetries are broken: localised turbulent patches of irregular shapes and various sizes which are surrounded by laminar regions can be observed for Re above about 320 (Bottin *et al.*, 1998*a,c*). The systems also allows for more ordered patterns of turbulent stripes which arise for a small range of parameters near $Re = 400$ and were reproduced in numerical simulations (Barkley & Tuckermann, 2005). For these Re localised perturbations are observed to generate localised turbulent spots that invade the surrounding laminar flow (Bottin *et al.*, 1998*b,c*).

As in previous studies in small periodic domains, we determine the edge state by numerically tracking the evolution of velocity fields which neither become fully turbulent nor decay to laminar flow but remain in regions intermediate between these two types of dynamics (Itano & Toh, 2001; Skufca *et al.*, 2006; Schneider *et al.*, 2007; Vollmer *et al.*, 2009). For the numerical simulation we use the Fourier-Chebyshev-tau scheme developed by Gibson (2004) with a resolution of 33 modes in normal direction. In the other directions, we adjust the number of modes when varying the size of the domain so that we keep $16/\pi$ modes per length in the spanwise and $4/\pi$ or $8/\pi$ modes per length in the downstream direction. One might expect that computing *localised* edge states requires a control not only on the perturbation energy but also on the spatial extension of a flow structure. However, as became clear in hindsight and will be demonstrated below, the evolution of these states is such that no additional control is needed and that the numerical algorithm described before (Schneider *et al.*, 2008; Schneider & Eckhardt, 2009) can be used without modification.

A domain that is 2π wide and 4π long suffices to support turbulent dynamics and is close to optimal for the appearance of coherent structures (Clever & Busse, 1997; Waleffe, 2003). We first focus on $Re = 400$, keep the length of the reference domain and extend its width to 8π and then 16π . In contrast to the case of the small domain, where a non-localised state has been found, the edge tracking algorithm now converges to a state that is localised in the spanwise direction, as shown in Fig. 1 (b). This state is not symmetric under reflection on the mid-plane and hence is not stationary but a travelling wave that moves




Figure 1: Symmetric (a), antisymmetric (b) and extended state (c) in plane Couette flow for $Re = 400$. Shown are downstream averages of the in-plane velocity components (arrows) and the downstream component (colour coded relative to the mean profile). (a) shows a fixed point symmetric with respect to a rotation around the centre. (b) shows a travelling wave invariant under the reflection $[u_x, u_y, u_z](x, y, z) \rightarrow [u_x, u_y, -u_z](x, y, -z)$. (c) shows the periodically continued edge state obtained from smaller domains. Note that the spanwise wavelength of the localised state is a bit shorter than the one for the extended state.




Figure 2: Properties of the edge state at $Re = 400$. The spectrum of the state (a) shows one unstable eigenvalue, two neutral related to the continuous translational symmetries and several stable real and complex ones. The energy of the edge state (b) quickly saturates as a function of the domain width L_z indicating its independence of the domain size.

with a phase speed of $6.9 \cdot 10^{-3} u_0$ downstream. There is a reflected partner travelling in the opposite direction. In the core region the state is dominated by pairs of downstream vortices that induce alternating high- and low-speed streaks. The topology is similar to the three-dimensional state described by (Nagata, 1990; Clever & Busse, 1997; Waleffe, 2003) and the non-localised edge state in small domains (Schneider *et al.*, 2008).

The eigenvalue spectrum shown in Fig. 2(a) confirms the conclusion drawn from the convergence of the edge state tracking, namely that the stable manifold is of co-dimension one. The variation of the total energy content with the box width shown in Fig. 2(b) confirms the localisation properties: the energy first increases but then settles to an essentially constant value once the width exceeds 7π . The full width at half maximum of the pattern this is about 8. This confirms that the properties are intrinsically controlled by the dynamics and not induced by the boundaries.

Using a Newton algorithm the travelling waves can be pinned and followed to different Reynolds numbers. The kinetic energy in Fig. 3(a) and the spanwise size as determined from a Gaussian fit in Fig. 3(b) show that the solution is spatially extended for low Re , localises as Re increases and reaches a constant width beyond $Re \approx 250$. The state's phase velocity (Fig. 3(c)) first deviates from zero at $Re \approx 150$ and oscillates with Re . The next to leading eigenvalue is shown in Fig. 3(d): its real part becomes negative for Re slightly above 200 and confirms that the stable manifold of the travelling wave is of co-dimension one and that is an edge state.

The presence of two locally attracting travelling waves on the edge calls for an explanation of how the their stable manifolds and the two local boundaries between laminar and turbulent dynamics which they define can be sewed together. The simplest explanation (following the bifurcation scenarios discussed




Figure 3: Properties of edge state for different Re . The variation of energy (a) and of the width (b) of the states with Re shows rather little variation above about 250. The phase velocity of the states (c) shows oscillations, and vanishes near about $Re = 220$ and 480. Both curves correspond to the two symmetry related solutions travelling in up- and down-stream direction. The next to leading eigenvalue (d) becomes negative for Re above 200 only. The interpolating lines are added to guide the eye only.




Figure 4: Bifurcation diagram for the localised travelling wave and the stationary state. The projection is defined by $\Sigma = \langle \vec{u}^2 \rangle_{x,y,z} - 7.5 \langle |\langle \vec{u}^2 \rangle_{x,z}(y) - \langle \vec{u}^2 \rangle_{x,z}(-y)| \rangle_y$. Insets: z -dependent spanwise velocity averaged in downstream and wall-normal direction.

in (Vollmer *et al.*, 2009)) requires the existence of a relative saddle in the edge which has an unstable direction pointing to either of the states. The broken up-down symmetry of the travelling waves then suggests that such a state should be symmetric. A Newton search starting from a suitably tailored initial condition indeed converges to the symmetric and stationary state shown in Fig. 1 (a). Its eigenvalue spectrum shows the second unstable eigenvalue required to connect it to the travelling waves via a symmetry breaking bifurcation. Indeed, following both the travelling wave and the stationary state down to a Reynolds number close to 150.2, they merge. However, when approaching this point, the widths of the states increases (cf. Fig. 3), until they extend over the full domain near the bifurcation point. This is documented in Fig. 4, where we characterise the states using a specially tailored measure of the energy content. The quantity Σ is calculated from the energy difference between the state and its mirror image in the spanwise direction so that it amplifies the difference between the two localised states and shows the reduction in energy when they become localised. One notes that as the Reynolds number is reduced, all three solutions converge near $Re = 150.2$, showing that both the symmetric and antisymmetric localised state emerge out of the spatially extended equilibrium.

Similar localisation phenomena have been observed in homoclinic snaking scenarios (Knobloch, 2008; Burke & Knobloch, 2007; Dawes *et al.*, 2007). The similarities in phenomenology are remarkable, and become particularly clear when the flow variations in downstream direction are averaged out and only the spanwise velocity averaged in x and y is shown: According to the symmetries of the full 3-d velocity fields the averaged velocities come in patterns of either reflection or a point mirror symmetry, as shown in the insets in Fig. 4. Remarkably, the localised patterns in the 1-d Swift-Hohenberg model with cubic-quintic nonlinearity show the same symmetries. We also noticed that the states in the Swift-Hohenberg model and the ones obtained here can be scaled and superimposed to look almost identical: while such a quantitative agreement cannot be expected because of the different form of the equations from which they are obtained, it does underline the strong similarities between the two systems, thereby pointing to a similar localisation mechanism.

3 Edge states in wide and long domains

Turning to domains that are 2π wide but much longer than 4π we find edge states that are localised in the downstream direction. However, their length falls off rather slowly, so that for $Re = 400$, where the length is about 60 to 80, boxes of a length 64π had to be used before localisation could be seen. As in the wide box this edge state is dominated by streaks but it is neither a fixed point nor a travelling wave but shows constant internal dynamics similar to the chaotic edge state found in pipe flow (Schneider *et al.*, 2007). Also the localisation in downstream direction is not unlike the one seen in models and in full numerical simulations for pipe flow (Duguet *et al.*, 2008; Mellibovsky *et al.*, 2009; Marinc *et al.*, 2009).




Figure 5: Localised turbulence seed at $Re = 400$. (a) shows the streamwise velocity in the $y=0$ -plane, emphasising the streaky structures. The bottom frames highlight the localisation in downstream (b) and spanwise (c) directions, by showing the energy averaged over the transverse directions. Note the exponential localisation in downstream direction and the faster than exponential one in spanwise direction.

Increasing both the width and length of the computational domain to 128 times the area of the reference domain, the edge tracking algorithm converges to a structure that is fully localised in span- and streamwise direction. Fig. 5(a) shows the localised state for a domain with $L_x = 64\pi$ and $L_z = 16\pi$. Most of the energy density of the perturbation is concentrated within a length of 20 and a width of 5 as measured by the variance of the averaged kinetic energy distribution. The visual appearance including the tails of the structures is a bit larger, about 20×80 . The localised state shows a streaky structure and combines the localisation features observed in long but narrow and in short but wide domains: it is exponentially localised in streamwise direction (Fig. 5b) and super-exponentially in spanwise direction Fig. 5c). Data for different independent edge state calculations starting from different initial conditions and computed with varying domain sizes and differing numerical resolution has been included in those figures. The overlap of the data shows that both spatial extensions and energy distributions are dynamically selected and independent of the size of the computational domain, the numerical resolution and the initial condition.

The fully localised edge state is not stationary or a travelling wave but shows chaotic temporal and spatial variations. As for the edge states identified in short segments of pipe flow (Schneider & Eckhardt, 2006; Schneider *et al.*, 2007), the mild chaotic variations can be clearly distinguished from turbulence because of




Figure 6: Mean amplitude E_t , height and width characterised by the variances V_x and V_z of the averaged energy distribution of a spot near the edge of chaos which swings up to turbulence. Note that the spot does not grow in size until the growth in amplitude is nearly finished.

their limited variability in space and energy, and their fairly slow dynamics.

The significance of the localised edge state lies in their finite size which defines the length, width and topology of marginally self-sustained perturbations. They are the smallest self-sustained structures away from the laminar profile and are critical in the sense that weaker perturbations will decay and stronger ones will increase to become turbulent. In the full state space of the system it is their stable manifold that separates laminar from turbulent dynamics. Interestingly, the size of this edge state is also very close to the minimal spot size required to stimulate growth at constant front velocity determined experimentally in (Tillmark & Alfredsson, 1992).

The dynamical relevance of these localised edge states for the transition is further clarified in Fig. 6 where the evolution from an edge state into a turbulent spot is presented. As the flow becomes turbulent the time traces of the spatial extension (length and width) and of the energy density stored in the perturbation field reveal two stages of the transition process: First the energy increases while maintaining the size of the spot, and only once the interior has reached turbulence levels does it start to grow in width and length. Thus, the structure has to become turbulent locally before it can start to fill the domain. Incidentally, this property of the localised structures explains why they can be detected and determined by monitoring the energy only: the alternative path by which energy could increase, namely by spreading in space while keeping the local energy density of the edge state, does not happen.

4 Conclusions

We have computed the energetically minimal self-sustained perturbations in extended plane Couette flow and shown that they are spatially localised. These

states are naturally related to earlier turbulence transition studies in which localised perturbations of a fixed type were used to generate growing turbulent spots (Tillmark & Alfredsson, 1992). The experimental results in (Tillmark & Alfredsson, 1992) together with the numerical studies in (Lundbladh & v. Johansson, 1991) show that a perturbation has to exceed a critical amplitude in order to generate a constantly growing spot. Remarkably, these studies also suggest that the critical perturbations are dominated by downstream vortices of size and topology very similar to the edge state shown in Fig. 5. This further supports the significance of localised edge states as nuclei for the transition dynamics.

The internal dynamics of the localised edge states can be complicated: in a wide but short domain they are travelling waves (cf. Fig. 1) of a topology similar to the non-localised edge states found in small periodically continued domains. In a wide and long domain the critical state is temporally active but shows a very limited spatial complexity when compared to a turbulent flow field. These observations together with the 2-d map studied in Vollmer *et al.* (2009) suggest that the relative attractor in the edge can be as simple or as complicated as a regular attractor in the full state space.

The localization properties discussed here introduce a new length scale to the system: comparing the observations of localised critical structures in plane Couette flow and pipe flow one notes that in both cases the structures have a localisation length that is much larger than the intrinsic structures of the edge states in the small domain, but that is shorter than the diameter of turbulent patches. In principle, models of front dynamics (e.g. Schumacher & Eckhardt (2001) and references therein) could help here, but their derivation starting from the Navier-Stokes equation remains a challenge.

Aspects of the spatial evolution of turbulent patches in spatially extended systems have been considered by Pomeau (1986), who suggested that the transition could have similarities to nucleation phenomena in first order equilibrium phase transitions. It is well known that in such cases a sufficiently strong perturbation is needed to induce the transition from one phase to the other. For instance, water droplets in a saturated water vapour dissolve if they are too small, and grow rapidly once they are sufficiently big. The same behaviour can be observed in the localised structures discussed here: if they are too weak or too small, they decay, and only if they exceed the relevant thresholds do they increase and spread. Pomeau (1986) proposed that there should be an appropriate non-equilibrium generalisation of the equilibrium phase-transition problem. The localised edge states shown here seem to be this non-equilibrium equivalent of the critical size droplets, and could be important for other aspects of the spatio-temporal dynamics in large domain turbulence as well Manneville (2009).

The work presented here is based on the diploma thesis of Daniel Marinc, completed in June 2008. Parts of the results were previously presented at the Newton Institute "Workshop on Wall bounded shear flows", Cambridge, Sept 8-12, 2008, and the 7th ERCOFTAC SIG33 Workshop "Open issues in transition and flow control", Genua, October 16-18, 2009. We thank the participants

of these meetings for discussion and the Deutsche Forschungsgemeinschaft for support.

References

- BARKLEY, DWIGHT & TUCKERMANN, LAURETTE S. 2005 Computational study of turbulent laminar patterns in Couette flow. *Physical Review Letters* **94**, 014502.
- BOBERG, L. & BROSA, U. 1988 Onset of turbulence in a pipe. *Z. Naturforsch.* **43a**, 697–726.
- BOTTIN, SABINE, DAUCHOT, OLIVIER & DAVIAUD, FRANCOIS 1997 Intermittency in a locally forced plane Couette flow. *Physical Review Letters* **79**, 4377–4380.
- BOTTIN, SABINE, DAUCHOT, OLIVIER, DAVIAUD, FRANCOIS & MANNEVILLE, PAUL 1998*a* Experimental evidence of streamwise vortices as finite amplitude solutions in transitional plane Couette flow. *Physics of Fluids* **10**, 2597–2607.
- BOTTIN, SABINE, DAVIAUD, FRANCOIS, MANNEVILLE, PAUL & DAUCHOT, OLIVIER 1998*b* Discontinuous transition to spatiotemporal intermittency in plane Couette flow. *Europhysics Letters* **43**, 171–176.
- BOTTIN, S., DAVIAUD, F., MANNEVILLE, P. & DAUCHOT, O. 1998*c* Discontinuous transition to spatiotemporal intermittency in plane couette flow. *Europhys. Lett.* **43** (2), 171 – 176.
- BURKE, JOHN & KNOBLOCH, EDGAR 2007 Homoclinic snaking: Structure and stability. *Chaos* **17**, **037102**, 15 pages.
- CLEVER, R.M. & BUSSE, FRITZ H. 1997 Tertiary and quaternary solutions for plane Couette flow. *Journal of Fluid Mechanics* **344**, 137–153.
- DARBYSHIRE, A.G. & MULLIN, T. 1995 Transition to turbulence in constant-mass-flux pipe flow. *J. Fluid Mech.* **289**, 83–114.
- DAUCHOT, OLIVIER & DAVIAUD, FRANCOIS 1995 Finite amplitude perturbation and spots growth mechanism in plane Couette flow. *Physics of Fluids* **7**, 335–343.
- DAWES, J.H.P., BURKE, J., CHAPMAN, S.J., KNOBLOCH, E., KOZYREFF, G., LLOYD, D. & MATTHEWS, P.C. 2007 Localized pattern formation p. 11 pages.
- DUGUET, Y., WILLIS, A. P. & KERSWELL, R. R. 2008 Transition in pipe flow: the saddle structure on the boundary of turbulence. *Journal of Fluid Mechanics* **???**, **???**, j. Fluid Mech., 2007, 21 pages.

- ECKHARDT, BRUNO, FAISST, HOLGER, SCHMIEGEL, ARMIN & SCHNEIDER, TOBIAS M 2008 Dynamical systems and the transition to turbulence in linearly stable shear flows. *Philosophical Transactions Royal Society London A* **366**, 1297–1315.
- ECKHARDT, BRUNO, FAISST, HOLGER, SCHMIEGEL, ARMIN & SCHUMACHER, JÖRG 2002 Turbulence transition in shear flows. In *Advances in Turbulence IX* (ed. I P Castro, P E Hancock & T G Thomas), pp. 701–708. CIMNE (Barcelona).
- ECKHARDT, BRUNO, SCHNEIDER, TOBIAS M, HOF, BJORN & WESTERWEEL, JERRY 2007 Turbulence transition in pipe flow. *Annual Review of Fluid Mechanics* **39**, 447–468.
- EMMONS, H W 1951 The laminar-turbulent transition in a boundary layer. *Journal Aeronautical Science* **18**, 490–498.
- GIBSON, JOHN F 2004 www.channelflow.org. *Tech. Rep.*.
- GROSSMANN, S 2000 The onset of shear flow turbulence. *Rev. Mod. Phys.* **72** (2), 603–618.
- HOF, B, JUEL, A & MULLIN, T 2003 Scaling of the turbulence transition threshold in a pipe. *Physical Review Letters* **91**, 244502.
- HOF, BJÖRN, WESTERWEEL, JERRY, SCHNEIDER, TOBIAS M & ECKHARDT, BRUNO 2006 Finite lifetime of turbulence in shear flows. *Nature* **443**, 60–64.
- ITANO, TOMOAKI & TOH, SADAYOSHI 2001 The dynamics of bursting process in wall turbulence. *Journal Physical Society Japan* **70**, 703–716.
- KNOBLOCH, E. 2008 Spatially localized structures in dissipative systems: open problems. *Nonlinearity* **21**, T45–T60 (16 pages).
- KOSCHMIEDER, E. L. 1993 *Bénard Cells and Taylor Vortices*. Cambridge University Press.
- LANDAU, L.D. 1944 On the problem of turbulence. *C.R. Acad. Sci. USSR* **44**, 311–314.
- LUNDBLADH, ANDERS & V. JOHANSSON, ARNE 1991 Direct simulation of turbulent spots in plane couette flow. *J. Fluid Mech.* **229**, 499–516 (18 pages).
- MANNEVILLE, PAUL 2009 Spatiotemporal perspective on the decay of turbulence in wall-bounded flows. *Physical Review E* **79**, 025301(R).
- MARINC, DANIEL, SCHNEIDER, TOBAS M & ECKHARDT, BRUNO 2009 Localise edge states for the transition to turbulence in shear flows. In *IUTAM Symposium Shear Flows and turbulence transition* (ed. D.S. Henningson & P. Schlatter).

- MELIBOVSKY, FERNANDO, MESEGUER, ALVARO, SCHNEIDER, TOBIAS M & ECKHARDT, BRUNO 2009 Transition in localized pipe flow turbulence. *Physical Review Letters* **103**, 054502.
- MESEGUER, ALVARO 2003 Streak breakdown instability in pipe poiseuille flow. *Physics of Fluids* **15**, 1203–1213.
- NAGATA, M 1990 Three-dimensional finite-amplitude solutions in plane Couette flow : bifurcation from infinity. *Journal of Fluid Mechanics* **217**, 519–527.
- PEIXINHO, JORGE & MULLIN, TOM 2007 Finite-amplitude thresholds for transition in pipe flow. *Journal of Fluid Mechanics* **582**, 169–178.
- POMEAU, Y. 1986 Front motion, metastability and subcritical bifurcations in hydrodynamics. *Physica D* **23**, 3–11 (9 pages).
- REYNOLDS, OSBORNE 1883 An experimental investigation of the circumstances which determine whether the motion of water shall be direct or sinuous, and the law of resistance in parallel channels. *Philosophical Transactions Royal Society (London)* **174**, 935–982 + 3 plates.
- SCHMID, P. J. & HENNINGSON, D. S. 1999 *Stability and Transition of Shear Flows*. New York: Springer.
- SCHMIEGEL, ARMIN & ECKHARDT, BRUNO 1997 Fractal stability border in plane couette flow. *Physical Review Letters* **79**, 5250–5253.
- SCHNEIDER, TOBIAS M & ECKHARDT, BRUNO 2006 Edge of chaos in pipe flow. *Chaos* **16**, 041103 (1 page).
- SCHNEIDER, TOBIAS M & ECKHARDT, BRUNO 2009 Edge states intermediate between laminar and turbulent dynamics in pipe flow. *Philosophical Transactions Royal Society London A* **367**, 577–587.
- SCHNEIDER, TOBIAS M, ECKHARDT, BRUNO & YORKE, JAMES A 2007 Turbulence transition and the edge of chaos in pipe flow. *Physical Review Letters* **99**, 034502.
- SCHNEIDER, TOBIAS M, GIBSON, JOHN F, LAGHA, MAHER, DELILLO, FILIPPO & ECKHARDT, BRUNO 2008 Laminar-turbulent boundary in plane couette flow. *Physical Review E* **78**, 037301 (4 pages).
- SCHUMACHER, JÖRG & ECKHARDT, BRUNO 2001 Evolution of turbulent spots in a parallel shear flow. *Physical Review E* **63**, 046307.
- SKUFCA, JOSEPH D, YORKE, JAMES A & ECKHARDT, BRUNO 2006 Edge of chaos in a parallel shear flow. *Physical Review Letters* **96**, 174101 (4 pages).
- TILLMARK, NILS & ALFREDSSON, P HENRIK 1992 Experiments on transition in plane couette flow. *Journal of Fluid Mechanics* **235**, 89–102.

- VOLLMER, JÜRGEN, SCHNEIDER, TOBIAS M & ECKHARDT, BRUNO 2009 Basin boundary, edge of chaos, and edge state in a two-dimensional model. *New Journal of Physics* **11**, 013040 (23pp).
- WALEFFE, FABIAN 2003 Homotopy of exact coherent structures in plane shear flows. *Physics of Fluids* **15**, 1517–1534.
- WANG, JUE, GIBSON, JOHN & WALEFFE, FABIAN 2007 Lower branch coherent states in shear flows: transition and control. *Physical Review Letters* **98**, 204501.